

# Diabetes Medical Management Plan

School District: \_\_\_\_\_ School: \_\_\_\_\_ School Year: \_\_\_\_\_ Grade: \_\_\_\_\_

Student Name: \_\_\_\_\_ DOB: \_\_\_\_\_

Provider Name: \_\_\_\_\_ Phone #: \_\_\_\_\_ Fax #: \_\_\_\_\_

## Blood Glucose Monitoring at School

Blood Glucose Target Range: \_\_\_\_\_ - \_\_\_\_\_ mg/dl

### Monitoring Schedule:

- Before breakfast     Before lunch     10-20 min. before boarding bus     Suspected hyper/hypoglycemia
- Is ill or requests testing     Other: \_\_\_\_\_

### Student Self Monitoring (Check all that apply.):

- Can test independently     Needs supervision     Needs assistance with testing and blood glucose management
- Other: \_\_\_\_\_

## Diabetes Medication

Oral medications: Home: \_\_\_\_\_ School: \_\_\_\_\_

Insulin: (*Opened insulin must be discarded after 28 days.*)

- No insulin at School    Insulin at Home:  Humalog     Novolog     Lantus Other: \_\_\_\_\_
- Insulin at School:  Humalog     Novolog     Lantus Other: \_\_\_\_\_

### Insulin delivery devise at school:

- Syringe & vial     Insulin Pen     Insulin Pump (See Pump Section.)

### Insulin management at school:

- |            |  |   |   |
|------------|--|---|---|
| Student is | Give own injections.                                 | <input type="checkbox"/> Y <input type="checkbox"/> N | <input type="checkbox"/> With supervision |
| able to:   | Draw up correct dose of insulin.                     | <input type="checkbox"/> Y <input type="checkbox"/> N | <input type="checkbox"/> With supervision |
|            | Determine correct amount of insulin.                 | <input type="checkbox"/> Y <input type="checkbox"/> N | <input type="checkbox"/> With supervision |
|            | Independently self manage pump or insulin injection. | <input type="checkbox"/> Y <input type="checkbox"/> N | <input type="checkbox"/> With supervision |

## Meals & Snacks at School

Independent in Carbohydrate calculations and management:     Yes     No     Needs Supervision

Meal/Snack	Carbohydrate Count	Not on Fixed Carb Count	Meal/Snack	Carbohydrate Count	Not on Fixed Carb Count
Breakfast			Lunch		
Mid-morning Snack			Mid-morning Snack		

Snack before exercise:     Yes     No     As Needed    Snack after exercise:     Yes     No     As Needed

Snack/content/amount at other times:     As Needed    OR    \_\_\_\_\_

Food to avoid: Liquid sugars such as fruit juice, regular soda and Gatorade. Use only for low blood sugars.  
Other: \_\_\_\_\_

Instructions when food provided in classroom (e.g. class party, food sampling): \_\_\_\_\_



### Hypoglycemia (Low Blood Sugar) = \_\_\_\_\_mg/dl and/or Physical Symptoms

**Symptoms of Hypoglycemia:**

Shaky      Headache      Confused      Clumsy      Sweaty      Drowsy      Hungry      Pale  
Uncooperative      Irritable      Weak      Behavior Changes      Other: \_\_\_\_\_

**Precautions**

- **Never leave this student unattended! If treatment is to be provided in the Health Office, a responsible adult needs to accompany the student to the Health Office.**
- **Check blood sugar if student has not done so and is symptomatic.**
- **Notify School Nurse and Parent when any of the following treatments are performed.**

**Low Blood Sugar Treatment:**

- Give ½ cup (4 oz.) of juice or regular soda or 3-4 glucose tablets (or 15 grams of fast acting carbohydrate). Do not cover with insulin. The carbohydrate is given to treat the low blood sugar.
- Recheck blood glucose in 15 minutes. If blood sugar is still **below** \_\_\_\_\_ give another 15 grams of carbohydrate.
- If the student's blood sugar is **above** \_\_\_\_\_, give a 15-30 gram carbohydrate snack or lunch.
- Make sure the student feels well before sending to lunch.
- Comments \_\_\_\_\_

**Treatment if disoriented, combative, and incoherent but is conscious:**

- Give ½ to 1 tube of glucose gel or cake decorating gel. Place gel between cheek and gum.
- Massage the outside of cheek to facilitate absorption through the membrane of the cheek.
- Encourage student to swallow.
- Recheck blood sugar in 10 minutes.
- If still **below** \_\_\_\_\_, repeat treatment as above.
- Give sugar containing liquid and snack when student is alert and able to swallow safely.
- Comments \_\_\_\_\_

**Treatment for seizures, loss of consciousness, inability/unwillingness to take gel or juice:**

- Stay with student
- Position student on side
- Give glucagon immediately by injection. Dose:  0.3cc     0.5cc     1.0cc
- Call 911
- **Notify parents**
- Comments \_\_\_\_\_

### Hyperglycemia (High Blood Sugar) = 250 or 300 mg/dl

**Symptoms of Hyperglycemia:**

Extreme Thirst      Frequent Urination      Abdominal Pain      Headache      Nausea  
Other: \_\_\_\_\_

**Check Ketones:**

- Urine should be checked for ketones when blood glucose levels are above 300 mg/dl.
- If urine ketones are moderate to large, **CALL PARENT IMMEDIATELY!**
- If student is on pump, and urine ketones are moderate to large OR blood ketones are 0.6mmol/l or more, call parents.

**Treatment for ketones and/or high blood sugar:**

- Increase sugar free liquid intake
- Allow student to use restroom as often as necessary
- Call parent immediately if student is vomiting

**Treatment for high glucose with ketones, moderate, large or ≥ 0.6 or greater: (check all that apply)**

- Call parent immediately for action plan
- Parent will determine the insulin coverage needed
- Follow blood sugar correction guidelines – see dosing sheet

### Supplies Kept at School

- Blood glucose meter, test strips, meter batteries
- Meter location: \_\_\_\_\_
- Insulin, pen, pen needles, insulin cartridges
- Lancet device, lancets, gloves, etc.
- Glucagon Emergency Kit
- Urine ketone strips
- Insulin vials and syringes
- Carbohydrate containing snack
- Fast-acting source of glucose
- Blood ketone meter and strips
- Insulin pump and supplies

### Insulin Pump

Insulin Pump Care Information Attached

Student able to operate insulin pump:  Y  N  With Supervision

Student can troubleshoot problems:  Y  N  With Supervision  
(e.g. Urine Ketones, pump malfunction)

Comments: \_\_\_\_\_

\_\_\_\_\_

### Insulin Adjustments by Healthcare Provider or Parent (for use by School Nurse)

Date New Orders Obtained	Order * Note Change in Care Sheet	Nurse Signature
	<input type="checkbox"/> Verbal <input type="checkbox"/> Written	
	<input type="checkbox"/> Verbal <input type="checkbox"/> Written	
	<input type="checkbox"/> Verbal <input type="checkbox"/> Written	
	<input type="checkbox"/> Verbal <input type="checkbox"/> Written	

### SIGNATURES:

This Diabetes Medical Management Plan has been approved by:

\_\_\_\_\_  
Student Healthcare Provider      Phone      Date      E-mail

\_\_\_\_\_  
Diabetes Educator      Phone      Date      E-mail

I give my permission to the school, school nurse, licensed/unlicensed assistive personnel, and other designated staff member(s) to perform and carry out the diabetes care tasks as outlined by this Diabetes Medical Management Plan for my child, \_\_\_\_\_, and I acknowledge that I have received a copy of the signed plan.

I also consent to the release of the information contained in this plan to all staff and other adults who have custodial care of my child and who may need to know this information to maintain my child's health and safety. I will notify extra-curricular staff about health plan and care to be given during after school activities. I give my permission for the school nurse to contact my child's healthcare provider(s) regarding the above condition.

\_\_\_\_\_  
Parent/Guardian      Phone      Date      E-mail

Acknowledged and received by:

\_\_\_\_\_  
School Nurse      Phone      Date      E-mail

Change in Care Management Plan

Student Name: \_\_\_\_\_ DOB: \_\_\_\_\_ New Order Date: \_\_\_\_\_

Carbohydrate Counting and Correction

Food: \_\_\_\_\_ units of Humalog/Novolog for every \_\_\_\_\_ grams of carbohydrate.
Blood Sugar: \_\_\_\_\_ units of Humalog/Novolog for every \_\_\_\_\_ mg/dl over \_\_\_\_\_ mg/dl.

- Corrections for blood sugar can be made every 3 hours if needed.
• Unless otherwise stated, cover all carbohydrates and snacks with insulin. Do not cover carbs used to treat low blood sugar.

PRE-MEAL Humalog/Novolog Doses

Table with 2 main columns: Blood Sugar Corrections and Food Carbohydrate. Each row contains 'Under' and 'to' values for units and grams.

Lantus dose is: \_\_\_\_\_ AM \_\_\_\_\_ at bedtime.
Bed time corrections: \_\_\_\_\_
At bed time correct blood sugar level to \_\_\_\_\_

Bedtime & 3:00 AM Correction

Table with 2 columns for 'Under' and 'to' values, each with 'Units' and 'Grams' sub-columns.

If blood sugar is less than \_\_\_\_\_ at bedtime, give \_\_\_\_\_ grams of carbohydrate + protein without Humalog/Novolog coverage for this snack.

- Change in Carb Counting and Blood Sugar correction per parent (if applicable).
Change in Carb Counting and Blood Sugar correction per provider (if applicable).
Additional changes to Initial Orders: \_\_\_\_\_

Signature

Printed Name

**Why are insulin pumpers at risk for ketoacidosis?**

Pumpers have no long-acting insulin (Lantus or Levemir) in their bodies. If the flow of insulin from the pump stops, the body will make ketones very quickly.

**What are the signs of high ketones?**

- Nausea
- Stomach cramps
- Vomiting
- Trouble breathing

Usually blood sugar level is high when there is a high number of ketones, but ketoacidosis can occur if the blood sugar is under 200. A person may think he/she has the stomach flu when, in fact, he/she is becoming ill from high ketones. Symptoms are exactly the same. If insulin is not given immediately, ketoacidosis will result.

Test urine or blood for ketones if the following symptoms are present. (Check expiration date on strips; if blood ketone strips are past expiration date, the machine will not read them.)

- Feeling sick or nauseated
- Blood sugar over 300
- Blood sugar over 250 two times in a row

**Follow directions below if ketones are present.****Less than 0.6 mmol/l Blood Ketones OR Trace/Small Urine Ketones**

- ADMINISTER correction bolus through insulin pump.
- RECHECK blood sugar and ketones in 1 hour.
- GIVE 4-8 oz. sugar free liquids by mouth every hour.
- **If blood sugar not improved in one hour, ADMINISTER insulin correction dose by syringe in amount equal to that recommended by the bolus wizard for the current blood sugar level.**
- REMOVE catheter and REPLACE insulin, cartridge, tubing and catheter.
- RECHECK blood sugar in two hours.
- ADMINISTER next bolus through pump with new set in place.

**0.6 mmol/l to 3.0 mmol/l Blood Ketones OR Moderate to Large Urine Ketones**

- ADMINISTER correction dose of fresh insulin by syringe **immediately** in amount equal to that recommended by bolus wizard for the current blood sugar level.
- GIVE 4-8 ounces sugar free liquids by mouth every hour.
- REMOVE catheter and REPLACE insulin, cartridge, tubing and catheter.
- RECHECK blood sugar and ketones every 2-3 hours.
- ADMINISTER next bolus through pump with new set in place.

**More than 3.0 mmol/l Blood Ketones**

- ADMINISTER **double** amount of correction insulin dose by syringe **immediately**.
- REMOVE catheter and REPLACE insulin, cartridge, tubing and catheter.
- CHECK blood sugar and ketones every 2-3 hours and set future correction doses using bolus wizard.
- ADMINISTER 4-8 oz. of sugar free liquids every hour.
- CALL the healthcare provider and parent/guardian.